arXiv:quant-ph/0103104v1 17 Mar 2001

Reply to Comment

Ping-Xing Chen and Lin-Mei Liang
Laboratory of Quantum Communication and Quantum Computation,
University of Science and Technology of China, Hefei, 230026, P. R. China
and
[*]*Department of Applied Physics,*
National University of Defense Technology, Changsha, 410073, P. R. China

Cheng-Zu Li
Department of Applied Physics,
National University of Defense Technology, Changsha, 410073, P. R. China

Ming-Qiu Huang
CCAST (World Laboratory) P.O. Box 8730, Beijing, 100080, China
and Department of Applied Physics,
National University of Defense Technology, Changsha, 410073, P. R. China

(February 1, 2008)

Typeset using REVTEX

---

[*] Corresponding address

In the comment [1] Eggeling, Vollbrecht and Wolf suspect our method in Ref. [2] is not practical. Here we explain our result and method and show that our example can tell one how to judge a separable state, and so our method is practical, at least for many mixed states.

A. Explanation

Recently we considered the necessary and sufficient condition of separability and gained a practical criterion [2]. Our main idea is as followings:

1. We decompose a mixed state ρ of $m \otimes n$ dimension Hilbert space into (m-1)(n-1) pairs which are in 2⊗2 dimension Hilbert space, from Ref. [3], we get the necessary condition of each pair being parallel, i.e. ρ being separable.

2. Each parallel pair corresponding to a set of matrices U, if there are intersection of all matrices U, ρ is separable, otherwise ρ is inseparable. We shown by our example [2] that judging the intersection existing or not comes down to judging a set of special symmetric quadratic equations having solutions or not, which is not difficult, so our method is practical.

B. Reply

The first step above generalizes Wootter's work [3] to higher dimension. This is meaningful not only for separable criterion but also for other approach(we are working in this). Parallel pair is different from PPT [4], though they are equivalent for 2⊗2 system, but in the case of higher dimension they have different character and usage.

If one has found a matrix U which brings up a set of separable pure states decomposition of ρ,one can say ρ is separable. But how to judge this matrix existing or not? The second step above has told one how to find this matrix U, so it is practical.

We think if the authors of the comment [1] consider the example in Ref [2] carefully, they will find our method is practical.

REFERENCES

[1] T.Eggeling, K.G.H.Vollbrecht, M.M.Wolf quant-ph/0103003

[2] P.X.Chen, Lin-Mei Liang, Cheng-Zu Li and Ming-Qiu Huang, quant-ph/0102133,Phys.Rev.A.(in press)

[3] W. K.Wootters, Phys. Rev. Lett. **80**, 2245(1998).

[4] A. Peres, Phys. Rev. Lett. **77**, 1413(1996).